



# OxiDate<sup>®</sup> 2.0

## PRODUCT BULLETIN

### FIFRA 2(ee) RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

### ISSUE DATE:

January 17, 2020

### EXPIRATION DATE:

June 1, 2023

### FOR USE AND DISTRIBUTION ONLY IN THE STATE OF:

**FLORIDA**

EPA REGISTRATION NO. 70299-12

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Crop:** Strawberries      **Diseases:** *Pestalotiopsis*

### Strawberry Application Instructions:

#### Pre-Plant Dip or Spray

Rate	Application	Notes
½-1.0 gallon of OxiDate 2.0 per 100 gallons of water.	Thoroughly wet transplants by dipping or spraying prior to planting.	Excessive foaming or bubbling during the dipping process is an indication of high levels of disease contamination.  Remove dead or dying foliage prior to dipping.

#### At-Planting Foliar Application

Rate	Application	Notes
1.0 gallon of OxiDate 2.0 per 100 gallons of water. Complete coverage is essential.	Immediately following planting, apply OxiDate 2.0 as a foliar spray with sufficient water to achieve runoff to soil or plastic, or to the soil directly via drip trickle, in furrow or flood basin.	Typical applications use 30-100 gallons of spray solution per treated acre, or 10-25 gallons of spray solution in low volume sprayers.

#### Existing Plantings Foliar Application

Rate – Foliar Spray	Application	Notes
<b>Preventative:</b> 1:200-1:400 or 64-32 fl. oz. per 100 gallons of water.	Begin preventative applications of OxiDate 2.0 prior to disease development and continue throughout the season, maintaining a 5-10 day spray schedule.	Typical applications use 30-100 gallons of spray solution per treated acre or 10-25 gallons of spray solution in low volume sprayers.
<b>Curative:</b> 1:100 or 1 gallon per 100 gallons of water.	Spray curative applications at first appearance of disease or when conditions are favorable for disease development. Repeat at 3 to 5 day intervals using sufficient water to obtain complete coverage.	Under severe disease conditions and during periods of rainy weather, apply immediately following each rain, reduce spray intervals and use 1:100 dilution rate.  Use sufficient water to obtain complete coverage.  OxiDate 2.0 may be applied up to and including the day of harvest.



BioSafe Systems, LLC

22 Meadow Street  
East Hartford, CT 06108

www.biosafesystems.com

1-888-273-3088

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use which are recommended by BioSafe Systems, LLC. which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This product bulletin must be in the possession of the user at the time of pesticide application.